

References Cited

American Geologic Institute. 1962. Dictionary of geologic terms. Garden City, NY: Doubleday and Company. 545 p.

Anderson, D and others. 1987. Idaho Department of Fish and Game, Regional Fishery Management Investigations, Job Performance Report Project F-71-R-11. Job No. 3 (MC)-c Subregion River and Stream Investigations. Rainbow trout fishery in the Cabarton reach of the North Fork Payette River. Pp 67-87.

Armantrout, NB, compiler. 1998. Glossary of aquatic habitat inventory terminology. Bethesda, MD: American Fisheries Society. 136 p.

Batt, PE. 1996. Governor Philip E. Batt's Idaho bull trout conservation plan. Boise, ID: State of Idaho, Office of the Governor. 20 p + appendices.

BCC. 1996. Gold Fork River Watershed Assessment Report. Boise, ID: Boise Cascade Corporation.

Beschta, RL. 1982. Comment on "Stream system evaluation with emphasis on spawning habitat for salmonids" by MA Shirazi and WK Seim. Water Resources Research 18:1292-1295.

Beschta, RL and Platts, WS. 1986. Morphological features of small streams: significance and function. American Water Resources Association, Water Resources Bulletin, Vol. 22(3): 369-379.

Bjornn, TC and Reiser, DW. 1991. Habitat requirements of salmonids in streams. American Fisheries Society Special Publication 19:83-138.

Borchardt, M.A. 1996. Nutrients. In Stevensen R.J., M.L. Bothwell, and R.L. Lowe editors. 1996. Algal Ecology: freshwater benthic ecosystems. San Diego CA: Academic Press, Inc.

Buening, JK. 2001. Water Quality Assessment for the Development of Total Maximum Daily Loads for Copper and Oil & Grease in Crawford Ditch in Elkhart County. Indianapolis, Indiana: Indiana Department of Environmental Management.

Carlson, RE. 1980. More complications in the chlorophyll-Secchi disc relationship. Limnology and Oceanography. 25:378-382.

Chapman DW and McLeod KP. 1987. Development of criteria for fine sediment in the Northern Rockies Ecoregion. Seattle, WA: Final Report to the Environmental Protection Agency. Clean Water Act (Federal water pollution control act), U.S.C. § 1251-1387 (1972).

CRWQCB. 2002. Mattole River Watershed, Technical Support Document for the Total Maximum Daily Loads for Sediment and Temperature. October 2002, Draft for Public Review. California Regional Water Quality Control Board North Coast Region.

Denny, P. 1980. Solute movement in submerged angiosperms. *Biology Review*. 55:65-92.

Dobson M and Drake M. 1990. Examining our Roots. Horseshoe Bend, ID: Scenic Payette River Historical Society. 96 p.

Dodds WK, Jones JR, Welch EB. 1998. Suggested classification of stream trophic state: distributions of temperate stream types by chlorophyll, total nitrogen, and phosphorus. *Water Research* 32: 1455-1462.

EPA. 1976. Quality Criteria for Water. Washington, DC: U.S. Environmental Protection Agency, (Red Book).

EPA, 1986. Quality Criteria for Water, 1986 (EPA 440/5-86-001) Washington, DC: U.S. Environmental Protection Agency, Office of Water. 162 p.

EPA. 1996. Biological Criteria: technical guidance for streams and small rivers. EPA 822-B-96-001. Washington, DC: U.S. Environmental Protection Agency, Office of Water. 162 p.

EPA. 1997. Guidelines for preparation of the comprehensive state water quality assessments (305(b) reports) and electronic updates: supplement. EPA-841-B-97-002B. Washington, DC: U.S. Environmental Protection Agency. 105 p.

EPA. 1999a. Protocol for Developing Nutrient TMDLs EPA 841-B-99-007. Washington DC, 20460: Office of Water 4503 F.

EPA. 1999b. Approval of EPA Methods 1664, Revision A, and 9071B for Determination of Oil and Grease and Non-polar Material in EPA's Wastewater and Hazardous Waste Programs. Washington, DC: U.S. Environmental Protection Agency, EPA-F-98-005.

EPA. 2000a. Ambient Water Quality Criteria Recommendations Information Supporting the Development of State and Tribal Nutrient Criteria Rivers and Streams in Nutrient Ecoregion II. Office of Water 4304, EPA 822-B-00-015.

EPA. 2000b. Ambient Water Quality Criteria Recommendations Information Supporting the Development of State and Tribal Nutrient Criteria Rivers and Streams in Nutrient Ecoregion III Office of Water 4304.0 EPA 822-B-00-016.

EPA. 2003. Decision Rationale Total Maximum Daily Loads Anacostia River Watershed and Kingman Lake for Oil and Grease. Philadelphia, PA: U.S. Environmental Protection Agency, Region III.

Ford BS, Higgins PS, Lewis AF, Cooper KL, Watson TA, Gee CM, Ennis GL and Sweeting RL. 1995. Literature reviews of the life history, habitat requirements and

mitigation/compensation strategies for thirteen sport fish species in the Peace, Liard and Columbia River drainages of British Columbia. Can. Manuscr. Rep. Fish. Aquat. Sci. 2321:xxiv+342p.

<http://www.georgiaconservancy.org/waterquality/>

Grafe CS, Mebane CA, McIntyre MJ, Essig DA, Brandt DH, Mosier DT. 2002. The Idaho Department of Environmental Quality water body assessment guidance, 2nd ed. Boise, ID: Department of Environmental Quality. 114 p.

Greenborg, AE, Clescevi LS, Eaton AD, editors. 1992. Standard methods for the examination of water and wastewater, 18th edition. Washington, DC: American Public Health Association. 900 p.

Hughes, RM. 1995. Defining acceptable biological status by comparing with reference condition. In: Davis WS, Simon TP, editors. Biological assessment and criteria: tools for water resource planning. Boca Raton, FL: CRC Press. p 31-48.

Idaho Code § 39.3611. Development and implementation of total maximum daily load or equivalent processes.

Idaho Code § 3615. Creation of watershed advisory groups.

Idaho Soil Conservation Commission. 2004. North Fork Payette River Clear Creek Round Valley Creet Riparian Assessment. David F. Ferguson, Idaho Soil Conservation Commission, and Guy Hopkins, Idaho Association of Soil Conservation Districts.

IDAPA 58.01.02. Idaho water quality standards and wastewater treatment requirements.

IDEQ. 1996. Cascade Reservoir Phase I Watershed Management Plan (Payette River Basin), Boise, ID: Idaho Department of Environmental Quality.

IDEQ. 1999. Lower Boise River Sediment and Bacteria Total Maximum Daily Load. Boise ID: Boise Regional Office, Idaho Department of Environmental Quality.

IDEQ. 2001. Upper Snake River-Rock Creek Subbasin Assessment and TMDL. Twin Falls, ID: Twin Falls Regional Office, Idaho Department of Environmental Quality.

IDEQ. 2002. Middle Salmon River-Chamberlain Creek Subbasin Assessment and Crooked Creek TMDL. Prepared by: Mark L. Shumar, Idaho Department of Environmental Quality, Boise.

IDEQ. 2003. Middle Snake River/Succor Creek Subbasin Assessment and Total Maximum Daily Load. Boise, ID: Boise Regional Office, Idaho Department of Environmental Quality.

IDEQ. 2003a. South Fork Clear Water Subbasin Assessment and Total Maximum Daily Loads. Lewiston, ID. Lewiston Regional Office, Idaho Department of Environmental Quality.

IDEQ. 2004. Snake River-Hells Canyon Subbasin Assessment and Total Maximum Daily Load. Boise, ID: Boise Regional Office, Idaho Department of Environmental Quality.

IDEQ. 2004a. South Fork Payette River Subbasin Assessment Boise, ID: Boise Regional Office, Idaho Department of Environmental Quality.

IDL. 2002. Big Creek Headwaters Cumulative Watershed Effects Assessment, CWE Assessment No. 17050123 – 1004. Coeur d'Alene, ID: Idaho Department of Lands.

IDL. 2003a. Box Creek Cumulative Watershed Effects Assessment, CWE Assessment No. 17050123 – 1107. Coeur d'Alene, ID: Idaho Department of Lands.

IDL. 2003b. Clear Creek Headwaters (HUC Nos. 17050123 – 1102) WQLS 2890. Cumulative Watershed Effects Assessment, CWE Assessment No. 17050123 – 1102. Coeur d'Alene, ID: Idaho Department of Lands.

IDL. 2003c. Clear Creek Sidewalls Cumulative Watershed Effects Assessment, CWE Assessment No. 17050123 – 1110. Coeur d'Alene, ID: Idaho Department of Lands.

IDL. 2003d. Fall Creek Cumulative Watershed Effects Assessment, CWE Assessment No. 17050123 – 1108. Coeur d'Alene, ID: Idaho Department of Lands.

IDWR. 1999. Idaho Comprehensive State Water Plan Summary, Payette River Basin. Boise, ID.

Irwin RJ, Van Mouwerik M, Stevens L, Seese MD, Basham W. 1997. Oil and Grease Entry. Environmental Contaminants Encyclopedia. Fort Collins, CO: National Park Service, Water Resources Division.

Karr JR. 1991. Biological integrity: a long-neglected aspect of water resource management. Ecological Applications. 1:66-84.

Kimmel, BL and Groeger AW. 1984. Factors controlling primary production in lakes and reservoirs, p. 277-281. IN Lake and reservoir management. Proc. North American Lake Management Society Symposium. EPA 440/5/84-001. Washington, DC: US EPA. 604 pp.

Lee GF, Jones-Lee A, Rast W. Alternative Approaches for Trophic State Classification for Water Quality Management, Part I: Suitability of Existing Trophic State Classification Systems, Part II: Application of Vollenweider-OECD Eutrophication Modeling Approach. El Macero, CA: Reports of G. Fred Lee & Associates. 1995.

MacDonald A. and Keller EA. 1987. Stream channel response to the removal of large woody debris, Larry Damm Creek, northwestern California. In: Erosion and sedimentation in the Pacific Rim. IAHS Publ. No. 165. Pages 405-406.

MacDonald LH, Smart AW, Wissmar RC. 1991. Monitoring guidelines to evaluate the effects of forestry activities on streams in the Pacific and Alaska. EPA/910/9-91-001. Seattle, WA: U.S. Environmental Protection Agency and University of Washington. 166 pp.

Megahan WF and Nowlin RA. 1976. *Sediment storage in channels draining small forested watersheds in the mountains of central Idaho*. Denver, CO: Third Federal Inter-agency Sedimentation Conference.

Newcombe CP and Jensen JOT. 1996. Channel suspended sediment and fisheries: a synthesis for quantitative assessment of risk and impact. *North American Journal of Fisheries Management*. 16:693-727.

NRCS. 1983. Proceedings from Natural Resource Conservation Service Channel Evaluation Workshop.

ODEQ. 2004a Willamette Basin TMDL: Appendix C, Potential Near-stream Land Cover in the Willamette Basin for Temperature Total Maximum Daily Loads (TMDLs). January 2004. Water Quality Division, Oregon Department of Environmental Quality.

ODEQ. 2004b. Walla Walla Subbasin Stream Temperature Total Maximum Daily Load and Water Quality Management Plan. November 19, 2004 Review Draft. Oregon Department of Environmental Quality.

Overton CK, McIntyre JD, Armstrong R, Whitwell S, Duncan KA. 1995. User's guide to fish habitat: natural conditions in the Salmon River Basin, Idaho. USDA Forest Service Technical Report INT-GTR-322. Ogden, UT: USDA Forest Service, Intermountain Research Station. 142 pp.

Platts WS, et al. 1987. Methods for evaluating riparian habitats with applications to management. USDA For. Serv. Gen. Tech. Rep. INT-221. 177 pp.

Platts WS, Meehan WF, Minshall GW. 1983. Methods for evaluating stream, riparian, and biotic conditions. General Technical Report INT-138. Ogden, UT: USDA Forest Service, Intermountain Forest and Range Experimental Station. 77 pp.

Rand GW, editor. 1995. Fundamentals of aquatic toxicology: effects, environmental fate, and risk assessment, second edition. Washington, DC: Taylor and Francis. 1125 p.

Rasmussen, LM. 1981. Soil Survey of Valley Area, Idaho, Parts of Adams and Valley Counties, USDS Soil Conservation Service in cooperation with the University of Idaho College of Agriculture, Idaho Soil Conservation Commission.

Rosgen DL. 1996. Applied River Morphology. Wildland Hydrology. Pagosa Springs, CO.

Schanz, F. and H. Juon. 1983. Two different methods of evaluating nutrient limitation of periphyton bioassays using water from the River Rhine and eight of its tributaries. *Hydrobiologia* 102, 187-195.

Strahler, AN. 1957. Quantitative analysis of watershed geomorphology. *American Geophysical Union Transactions*, 38:913-920.

Thornton, KW, Kimmel BL, Payne FE, editors. 1990. Reservoir Limnology: Ecological Perspective. New York, NY: John Wiley and Sons Inc. 253 p.

Thurston RV, Russo RC, Fetterolf Jr CM, Edsall TA, and Barber Jr YM. 1979. A review of the EPA Red Book: Quality criteria for water. Bethesda, MD: American Fisheries Society. 313 p.

BOR. 2004. Resource Management Plan Black Canyon Reservoir and Montour Wildlife Management Area July 2004 U.S. Department of the Interior, Pacific Northwest Region, Snake River Area Office.

USDA. 1999. A procedure to estimate the response of aquatic systems to changes in phosphorus and nitrogen inputs. Portland, OR: National Water and Climate Center, Natural Resources Conservation Service.

USFS. 1999. Long Prong Project Final Environmental Impact Statement Boise National Forest Cascade Ranger District. Chapter 4.

USFS 2003. BOISE National Forest Annual Summary Report 2002-2003 Effectiveness Monitoring Project for Streams and Riparian Areas within the Upper Columbia River Basin. PIBO Effectiveness Monitoring Project Staff.

USGS. 1987. Hydrologic unit maps. Denver, CO: United States Geological Survey. Water supply paper 2294. 63 p.

Water Pollution Control Federation. 1987. The Clean Water Act of 1987. Alexandria, VA: Water Pollution Control Federation. 318 p.

Water Quality Act of 1987, Public Law 100-4 (1987).

Water quality planning and management, 40 CFR 130.

Wetzel, R.T. 1975. Limnology. Philadelphia: Saunders College Publishing.

Wetzel, R.G. 1983. Limnology. Philadelphia: Saunders College Publishing. 793 pp.

Wetzel, R. 2001. Limnology, Lake and River Ecosystems. San Diego, CA: Academic Press.

WRCC (Western Regional Climate Center). March 10, 2004. Available from:
<http://www.wrcc.dri.edu/summary/climsmid.html>.

GIS Coverages:

Restriction of liability: Neither the state of Idaho nor the Department of Environmental Quality, nor any of their employees make any warranty, express or implied, or assume any legal liability or responsibility for the accuracy, completeness or usefulness of any information or data provided. Metadata is provided for all data sets, and no data should be used without first reading and understanding its limitations. The data could include technical inaccuracies or typographical errors. The Department of Environmental Quality may update, modify, or revise the data used at any time, without notice.

This page left blank intentionally.

Glossary

§305(b)	Refers to section 305 subsection “b” of the Clean Water Act. 305(b) generally describes a report of each state’s water quality, and is the principle means by which the U.S. Environmental Protection Agency, Congress, and the public evaluate whether U.S. waters meet water quality standards, the progress made in maintaining and restoring water quality, and the extent of the remaining problems.
§303(d)	Refers to Section 303 subsection “d” of the Clean Water Act. 303(d) requires states to develop a list of water bodies that do not meet water quality standards. This section also requires total maximum daily loads (TMDLs) be prepared for listed waters. Both the list and the TMDLs are subject to U.S. Environmental Protection Agency approval.
Acre-Foot	A volume of water that would cover an acre to a depth of one foot. Often used to quantify reservoir storage and the annual discharge of large rivers.
Adsorption	The adhesion of one substance to the surface of another. Clays, for example, can adsorb phosphorus and organic molecules
Aeration	A process by which water becomes charged with air directly from the atmosphere. Dissolved gases, such as oxygen, are then available for reactions in water.
Aerobic	Describes life, processes, or conditions that require the presence of oxygen.
Alevin	A newly hatched, incompletely developed fish (usually a salmonid) still in nest or inactive on the bottom of a water body, living off stored yolk.
Algae	Non-vascular (without water-conducting tissue) aquatic plants that occur as single cells, colonies, or filaments.
Alluvium	Unconsolidated recent stream deposition.
Ambient	General conditions in the environment. In the context of water quality, ambient waters are those representative of general conditions, not associated with episodic perturbations, or specific disturbances such as a wastewater outfall (Armantrout 1998, EPA 1996).
Anadromous	Fish, such as salmon and sea-run trout, that live part or the majority of their lives in the salt water but return to fresh water to spawn.

Anaerobic	Describes the processes that occur in the absence of molecular oxygen and describes the condition of water that is devoid of molecular oxygen.
Anoxia	The condition of oxygen absence or deficiency.
Anthropogenic	Relating to, or resulting from, the influence of human beings on nature.
Anti-Degradation	Refers to the U.S. Environmental Protection Agency's interpretation of the Clean Water Act goal that states and tribes maintain, as well as restore, water quality. This applies to waters that meet or are of higher water quality than required by state standards. State rules provide that the quality of those high quality waters may be lowered only to allow important social or economic development and only after adequate public participation (IDAPA 58.01.02.051). In all cases, the existing beneficial uses must be maintained. State rules further define lowered water quality to be 1) a measurable change, 2) a change adverse to a use, and 3) a change in a pollutant relevant to the water's uses (IDAPA 58.01.02.003.56).
Aquatic	Occurring, growing, or living in water.
Aquifer	An underground, water-bearing layer or stratum of permeable rock, sand, or gravel capable of yielding of water to wells or springs.
Assemblage (aquatic)	An association of interacting populations of organisms in a given water body; for example, a fish assemblage, or a benthic macroinvertebrate assemblage (also see Community) (EPA 1996).
Assimilative Capacity	The ability to process or dissipate pollutants without ill effect to beneficial uses.
Batholith	A large body of intrusive igneous rock that has more than 40 square miles of surface exposure and no known floor. A batholith usually consists of coarse-grained rocks such as granite.
Bedload	Material (generally sand-sized or larger sediment) that is carried along the streambed by rolling or bouncing.
Beneficial Use	Any of the various uses of water, including, but not limited to, aquatic biota, recreation, water supply, wildlife habitat, and aesthetics, which are recognized in water quality standards.
Beneficial Use Reconnaissance Program (BURP)	A program for conducting systematic biological and physical habitat surveys of water bodies in Idaho. BURP protocols address lakes, reservoirs, and Wadeable streams and rivers
Benthic	Pertaining to or living on or in the bottom sediments of a water body

Best Management Practices (BMPs)	Structural, nonstructural, and managerial techniques that are effective and practical means to control nonpoint source pollutants.
Biochemical Oxygen Demand (BOD)	The amount of dissolved oxygen used by organisms during the decomposition (respiration) of organic matter, expressed as mass of oxygen per volume of water, over some specified period of time.
Biological Integrity	1) The condition of an aquatic community inhabiting unimpaired water bodies of a specified habitat as measured by an evaluation of multiple attributes of the aquatic biota (EPA 1996). 2) The ability of an aquatic ecosystem to support and maintain a balanced, integrated, adaptive community of organisms having a species composition, diversity, and functional organization comparable to the natural habitats of a region (Karr 1991).
Biomass	The weight of biological matter. Standing crop is the amount of biomass (e.g., fish or algae) in a body of water at a given time. Often expressed as grams per square meter.
Biota	The animal and plant life of a given region.
Biotic	A term applied to the living components of an area.
Clean Water Act (CWA)	The Federal Water Pollution Control Act (commonly known as the Clean Water Act), as last reauthorized by the Water Quality Act of 1987, establishes a process for states to use to develop information on, and control the quality of, the nation's water resources.
Coliform Bacteria	A group of bacteria predominantly inhabiting the intestines of humans and animals but also found in soil. Coliform bacteria are commonly used as indicators of the possible presence of pathogenic organisms (also see Fecal Coliform Bacteria).
Community	A group of interacting organisms living together in a given place.
Conductivity	The ability of an aqueous solution to carry electric current, expressed in micro (μ) mhos/cm at 25 °C. Conductivity is affected by dissolved solids and is used as an indirect measure of total dissolved solids in a water sample.
Cretaceous	The final period of the Mesozoic era (after the Jurassic and before the Tertiary period of the Cenozoic era), thought to have covered the span of time between 135 and 65 million years ago.

Criteria	In the context of water quality, numeric or descriptive factors taken into account in setting standards for various pollutants. These factors are used to determine limits on allowable concentration levels, and to limit the number of violations per year. EPA develops criteria guidance; states establish criteria.
Cubic Feet per Second	A unit of measure for the rate of flow or discharge of water. One cubic foot per second is the rate of flow of a stream with a cross-section of one square foot flowing at a mean velocity of one foot per second. At a steady rate, once cubic foot per second is equal to 448.8 gallons per minute and 10,984 acre-feet per day.
Cultural Eutrophication	The process of eutrophication that has been accelerated by human-caused influences. Usually seen as an increase in nutrient loading (also see Eutrophication).
Decomposition	The breakdown of organic molecules (e.g., sugar) to inorganic molecules (e.g., carbon dioxide and water) through biological and nonbiological processes.
Designated Uses	Those water uses identified in state water quality standards that must be achieved and maintained as required under the Clean Water Act.
Discharge	The amount of water flowing in the stream channel at the time of measurement. Usually expressed as cubic feet per second (cfs).
Dissolved Oxygen (DO)	The oxygen dissolved in water. Adequate DO is vital to fish and other aquatic life.
Disturbance	Any event or series of events that disrupts ecosystem, community, or population structure and alters the physical environment.
<i>E. coli</i>	Short for <i>Escherichia Coli</i> , <i>E. coli</i> are a group of bacteria that are a subspecies of coliform bacteria. Most <i>E. coli</i> are essential to the healthy life of all warm-blooded animals, including humans. Their presence is often indicative of fecal contamination.
Ecology	The scientific study of relationships between organisms and their environment; also defined as the study of the structure and function of nature.
Ecological Indicator	A characteristic of an ecosystem that is related to, or derived from, a measure of a biotic or abiotic variable that can provide quantitative information on ecological structure and function. An indicator can contribute to a measure of integrity and sustainability. Ecological indicators are often used within the multimetric index framework.

Ecological Integrity	The condition of an unimpaired ecosystem as measured by combined chemical, physical (including habitat), and biological attributes (EPA 1996).
Ecosystem	The interacting system of a biological community and its non-living (abiotic) environmental surroundings.
Effluent	A discharge of untreated, partially treated, or treated wastewater into a receiving water body.
Endangered Species	Animals, birds, fish, plants, or other living organisms threatened with imminent extinction. Requirements for declaring a species as endangered are contained in the Endangered Species Act.
Environment	The complete range of external conditions, physical and biological, that affect a particular organism or community.
Ephemeral Stream	A stream or portion of a stream that flows only in direct response to precipitation. It receives little or no water from springs and no long continued supply from melting snow or other sources. Its channel is at all times above the water table. (American Geologic Institute 1962).
Erosion	The wearing away of areas of the earth's surface by water, wind, ice, and other forces.
Eutrophic	From Greek for "well nourished," this describes a highly productive body of water in which nutrients do not limit algal growth. It is typified by high algal densities and low clarity.
Eutrophication	1) Natural process of maturing (aging) in a body of water. 2) The natural and human-influenced process of enrichment with nutrients, especially nitrogen and phosphorus, leading to an increased production of organic matter.
Exceedance	A violation (according to DEQ policy) of the pollutant levels permitted by water quality criteria.
Existing Beneficial Use or Existing Use	A beneficial use actually attained in waters on or after November 28, 1975, whether or not the use is designated for the waters in Idaho's <i>Water Quality Standards and Wastewater Treatment Requirements</i> (IDAPA 58.01.02).
Exotic Species	A species that is not native (indigenous) to a region.
Extrapolation	Estimation of unknown values by extending or projecting from known values.
Fauna	Animal life, especially the animals characteristic of a region, period, or special environment.

Fecal Coliform Bacteria	Bacteria found in the intestinal tracts of all warm-blooded animals or mammals. Their presence in water is an indicator of pollution and possible contamination by pathogens (also see Coliform Bacteria).
Feedback Loop	In the context of watershed management planning, a feedback loop is a process that provides for tracking progress toward goals and revising actions according to that progress.
Fixed-Location Monitoring	Sampling or measuring environmental conditions continuously or repeatedly at the same location.
Flow	See Discharge.
Fluvial	In fisheries, this describes fish whose life history takes place entirely in streams but migrate to smaller streams for spawning.
Focal	Critical areas supporting a mosaic of high quality habitats that sustain a diverse or unusually productive complement of native species.
Fully Supporting	In compliance with water quality standards and within the range of biological reference conditions for all designated and existing beneficial uses as determined through the <i>Water Body Assessment Guidance</i> (Grafe et al. 2002).
Fully Supporting Cold Water	Reliable data indicate functioning, sustainable cold water biological assemblages (e.g., fish, macroinvertebrates, or algae), none of which have been modified significantly beyond the natural range of reference conditions (EPA 1997).
Geographical Information Systems (GIS)	A georeferenced database.
Geometric Mean	A back-transformed mean of the logarithmically transformed numbers often used to describe highly variable, right-skewed data (a few large values), such as bacterial data.
Grab Sample	A single sample collected at a particular time and place. It may represent the composition of the water in that water column.
Gradient	The slope of the land, water, or streambed surface.
Ground Water	Water found beneath the soil surface saturating the layer in which it is located. Most ground water originates as rainfall, is free to move under the influence of gravity, and usually emerges again as stream flow.
Growth Rate	A measure of how quickly something living will develop and grow, such as the amount of new plant or animal tissue produced per a given unit of time, or number of individuals added to a population.

Habitat	The living place of an organism or community.
Headwater	The origin or beginning of a stream.
Hydrologic Basin	The area of land drained by a river system, a reach of a river and its tributaries in that reach, a closed basin, or a group of streams forming a drainage area (also see Watershed).
Hydrologic Unit	One of a nested series of numbered and named watersheds arising from a national standardization of watershed delineation. The initial 1974 effort (USGS 1987) described four levels (region, subregion, accounting unit, cataloging unit) of watersheds throughout the United States. The fourth level is uniquely identified by an eight-digit code built of two-digit fields for each level in the classification. Originally termed a cataloging unit, fourth field hydrologic units have been more commonly called subbasins. Fifth and sixth field hydrologic units have since been delineated for much of the country and are known as watershed and subwatersheds, respectively.
Hydrologic Unit Code (HUC)	The number assigned to a hydrologic unit. Often used to refer to fourth field hydrologic units.
Hydrology	The science dealing with the properties, distribution, and circulation of water.
Impervious	Describes a surface, such as pavement, that water cannot penetrate.
Inorganic	Materials not derived from biological sources.
Instantaneous	A condition or measurement at a moment (instant) in time.
Intergravel Dissolved Oxygen	The concentration of dissolved oxygen within spawning gravel. Consideration for determining spawning gravel includes species, water depth, velocity, and substrate.
Intermittent Stream	1) A stream that flows only part of the year, such as when the ground water table is high or when the stream receives water from springs or from surface sources such as melting snow in mountainous areas. The stream ceases to flow above the streambed when losses from evaporation or seepage exceed the available stream flow. 2) A stream that has a period of zero flow for at least one week during most years.
Key Watershed	A watershed that has been designated in Idaho Governor Batt's <i>State of Idaho Bull Trout Conservation Plan</i> (1996) as critical to the long-term persistence of regionally important trout populations.

Land Application	A process or activity involving application of wastewater, surface water, or semi-liquid material to the land surface for the purpose of treatment, pollutant removal, or ground water recharge.
Limiting Factor	A chemical or physical condition that determines the growth potential of an organism. This can result in a complete inhibition of growth, but typically results in less than maximum growth rates.
Limnology	The scientific study of fresh water, especially the history, geology, biology, physics, and chemistry of lakes.
Load Allocation (LA)	A portion of a water body's load capacity for a given pollutant that is given to a particular nonpoint source (by class, type, or geographic area).
Load(ing)	The quantity of a substance entering a receiving stream, usually expressed in pounds or kilograms per day or tons per year. Loading is the product of flow (discharge) and concentration.
Loading Capacity (LC)	A determination of how much pollutant a water body can receive over a given period without causing violations of state water quality standards. Upon allocation to various sources, and a margin of safety, it becomes a total maximum daily load.
Loam	Refers to a soil with a texture resulting from a relative balance of sand, silt, and clay. This balance imparts many desirable characteristics for agricultural use.
Loess	A uniform wind-blown deposit of silty material. Silty soils are among the most highly erodible.
Luxury Consumption	A phenomenon in which sufficient nutrients are available in either the sediments or the water column of a water body, such that aquatic plants take up and store an abundance in excess of the plants' current needs.
Macroinvertebrate	An invertebrate animal (without a backbone) large enough to be seen without magnification and retained by a 500µm mesh (U.S. #30) screen.
Macrophytes	Rooted and floating vascular aquatic plants, commonly referred to as water weeds. These plants usually flower and bear seeds. Some forms, such as duckweed and coontail (<i>Ceratophyllum sp.</i>), are free-floating forms not rooted in sediment.

Margin of Safety (MOS)	An implicit or explicit portion of a water body's loading capacity set aside to allow the uncertainty about the relationship between the pollutant loads and the quality of the receiving water body. This is a required component of a total maximum daily load (TMDL) and is often incorporated into conservative assumptions used to develop the TMDL (generally within the calculations and/or models). The MOS is not allocated to any sources of pollution.
Mass Wasting	A general term for the down slope movement of soil and rock material under the direct influence of gravity.
Mean	Describes the central tendency of a set of numbers. The arithmetic mean (calculated by adding all items in a list, then dividing by the number of items) is the statistic most familiar to most people.
Median	The middle number in a sequence of numbers. If there are an even number of numbers, the median is the average of the two middle numbers. For example, 4 is the median of 1, 2, 4, 14, 16; and 6 is the median of 1, 2, 5, 7, 9, 11.
Metric	1) A discrete measure of something, such as an ecological indicator (e.g., number of distinct taxon). 2) The metric system of measurement.
Milligrams per liter (mg/L)	A unit of measure for concentration in water, essentially equivalent to parts per million (ppm).
Million gallons per day (MGD)	A unit of measure for the rate of discharge of water, often used to measure flow at wastewater treatment plants. One MGD is equal to 1.547 cubic feet per second.
Miocene	Of, relating to, or being an epoch of, the Tertiary between the Pliocene and the Oligocene periods, or the corresponding system of rocks.
Monitoring	A periodic or continuous measurement of the properties or conditions of some medium of interest, such as monitoring a water body.
Mouth	The location where flowing water enters into a larger water body.
National Pollution Discharge Elimination System (NPDES)	A national program established by the Clean Water Act for permitting point sources of pollution. Discharge of pollution from point sources is not allowed without a permit.
Natural Condition	A condition indistinguishable from that without human-caused disruptions.
Nitrogen	An element essential to plant growth, and thus is considered a nutrient.

Nonpoint Source	A dispersed source of pollutants, generated from a geographical area when pollutants are dissolved or suspended in runoff and then delivered into waters of the state. Nonpoint sources are without a discernable point or origin. They include, but are not limited to, irrigated and non-irrigated lands used for grazing, crop production, and silviculture; rural roads; construction and mining sites; log storage or rafting; and recreation sites.
Not Assessed (NA)	A concept and an assessment category describing water bodies that have been studied, but are missing critical information needed to complete an assessment.
Not Attainable	A concept and an assessment category describing water bodies that demonstrate characteristics that make it unlikely that a beneficial use can be attained (e.g., a stream that is dry but designated for salmonid spawning).
Not Fully Supporting	Not in compliance with water quality standards or not within the range of biological reference conditions for any beneficial use as determined through the <i>Water Body Assessment Guidance</i> (Grafe et al. 2002).
Not Fully Supporting Cold Water	At least one biological assemblage has been significantly modified beyond the natural range of its reference condition (EPA 1997).
Nuisance	Anything which is injurious to the public health or an obstruction to the free use, in the customary manner, of any waters of the state.
Nutrient	Any substance required by living things to grow. An element or its chemical forms essential to life, such as carbon, oxygen, nitrogen, and phosphorus. Commonly refers to those elements in short supply, such as nitrogen and phosphorus, which usually limit growth.
Nutrient Cycling	The flow of nutrients from one component of an ecosystem to another, as when macrophytes die and release nutrients that become available to algae (organic to inorganic phase and return).
Oligotrophic	The Greek term for “poorly nourished.” This describes a body of water in which productivity is low and nutrients are limiting to algal growth, as typified by low algal density and high clarity.
Organic Matter	Compounds manufactured by plants and animals that contain principally carbon.
Orthophosphate	A form of soluble inorganic phosphorus most readily used for algal growth.
Oxygen-Demanding Materials	Those materials, mainly organic matter, in a water body that consume oxygen during decomposition.

Parameter	A variable, measurable property whose value is a determinant of the characteristics of a system, such as temperature, dissolved oxygen, and fish populations are parameters of a stream or lake.
Partitioning	The sharing of limited resources by different races or species; use of different parts of the habitat, or the same habitat at different times. Also the separation of a chemical into two or more phases, such as partitioning of phosphorus between the water column and sediment.
Pathogens	Disease-producing organisms (e.g., bacteria, viruses, parasites).
Perennial Stream	A stream that flows year-around in most years.
Periphyton	Attached microflora (algae and diatoms) growing on the bottom of a water body or on submerged substrates, including larger plants.
Pesticide	Substances or mixtures of substances intended for preventing, destroying, repelling, or mitigating any pest. Also, any substance or mixture intended for use as a plant regulator, defoliant, or desiccant.
pH	The negative \log_{10} of the concentration of hydrogen ions, a measure which in water ranges from very acid (pH=1) to very alkaline (pH=14). A pH of 7 is neutral. Surface waters usually measure between pH 6 and 9.
Phased TMDL	A total maximum daily load (TMDL) that identifies interim load allocations and details further monitoring to gauge the success of management actions in achieving load reduction goals and the effect of actual load reductions on the water quality of a water body. Under a phased TMDL, a refinement of load allocations, wasteload allocations, and the margin of safety is planned at the outset.
Phosphorus	An element essential to plant growth, often in limited supply, and thus considered a nutrient.
Physiochemical	In the context of bioassessment, the term is commonly used to mean the physical and chemical factors of the water column that relate to aquatic biota. Examples in bioassessment usage include saturation of dissolved gases, temperature, pH, conductivity, dissolved or suspended solids, forms of nitrogen, and phosphorus. This term is used interchangeable with the terms “physical/chemical” and “physicochemical.”
Plankton	Microscopic algae (phytoplankton) and animals (zooplankton) that float freely in open water of lakes and oceans.

Point Source	A source of pollutants characterized by having a discrete conveyance, such as a pipe, ditch, or other identifiable “point” of discharge into a receiving water. Common point sources of pollution are industrial and municipal wastewater.
Pollutant	Generally, any substance introduced into the environment that adversely affects the usefulness of a resource or the health of humans, animals, or ecosystems.
Pollution	A very broad concept that encompasses human-caused changes in the environment which alter the functioning of natural processes and produce undesirable environmental and health effects. This includes human-induced alteration of the physical, biological, chemical, and radiological integrity of water and other media.
Population	A group of interbreeding organisms occupying a particular space; the number of humans or other living creatures in a designated area.
Pretreatment	The reduction in the amount of pollutants, elimination of certain pollutants, or alteration of the nature of pollutant properties in wastewater prior to, or in lieu of, discharging or otherwise introducing such wastewater into a publicly owned wastewater treatment plant.
Primary Productivity	The rate at which algae and macrophytes fix carbon dioxide using light energy. Commonly measured as milligrams of carbon per square meter per hour.
Protocol	A series of formal steps for conducting a test or survey.
Qualitative	Descriptive of kind, type, or direction.
Quantitative	Descriptive of size, magnitude, or degree.
Reach	A stream section with fairly homogenous physical characteristics.
Reconnaissance	An exploratory or preliminary survey of an area.
Reference	A physical or chemical quantity whose value is known, and thus is used to calibrate or standardize instruments.

Reference Condition	1) A condition that fully supports applicable beneficial uses with little effect from human activity and represents the highest level of support attainable. 2) A benchmark for populations of aquatic ecosystems used to describe desired conditions in a biological assessment and acceptable or unacceptable departures from them. The reference condition can be determined through examining regional reference sites, historical conditions, quantitative models, and expert judgment (Hughes 1995).
Reference Site	A specific locality on a water body that is minimally impaired and is representative of reference conditions for similar water bodies.
Representative Sample	A portion of material or water that is as similar in content and consistency as possible to that in the larger body of material or water being sampled.
Resident	A term that describes fish that do not migrate.
Respiration	A process by which organic matter is oxidized by organisms, including plants, animals, and bacteria. The process converts organic matter to energy, carbon dioxide, water, and lesser constituents.
Riffle	A relatively shallow, gravelly area of a streambed with a locally fast current, recognized by surface chopiness. Also an area of higher streambed gradient and roughness.
Riparian	Associated with aquatic (stream, river, lake) habitats. Living or located on the bank of a water body.
Riparian Habitat Conservation Area (RHCA)	A U.S. Forest Service description of land within the following number of feet up-slope of each of the banks of streams: <ul style="list-style-type: none"> - 300 feet from perennial fish-bearing streams - 150 feet from perennial non-fish-bearing streams - 100 feet from intermittent streams, wetlands, and ponds in priority watersheds.
River	A large, natural, or human-modified stream that flows in a defined course or channel, or a series of diverging and converging channels.
Runoff	The portion of rainfall, melted snow, or irrigation water that flows across the surface, through shallow underground zones (interflow), and through ground water to creates streams.
Sediments	Deposits of fragmented materials from weathered rocks and organic material that were suspended in, transported by, and eventually deposited by water or air.

Settleable Solids	The volume of material that settles out of one liter of water in one hour.
Species	1) A reproductively isolated aggregate of interbreeding organisms having common attributes and usually designated by a common name. 2) An organism belonging to such a category.
Spring	Ground water seeping out of the earth where the water table intersects the ground surface.
Stratification	A Department of Environmental Quality classification method used to characterize comparable units (also called classes or strata).
Stream	A natural water course containing flowing water, at least part of the year. Together with dissolved and suspended materials, a stream normally supports communities of plants and animals within the channel and the riparian vegetation zone.
Stream Order	Hierarchical ordering of streams based on the degree of branching. A first-order stream is an unforked or unbranched stream. Under Strahler's (1957) system, higher order streams result from the joining of two streams of the same order.
Storm Water Runoff	Rainfall that quickly runs off the land after a storm. In developed watersheds the water flows off roofs and pavement into storm drains that may feed quickly and directly into the stream. The water often carries pollutants picked up from these surfaces.
Subbasin	A large watershed of several hundred thousand acres. This is the name commonly given to 4 th field hydrologic units (also see Hydrologic Unit).
Subbasin Assessment (SBA)	A watershed-based problem assessment that is the first step in developing a total maximum daily load in Idaho.
Subwatershed	A smaller watershed area delineated within a larger watershed, often for purposes of describing and managing localized conditions. Also proposed for adoption as the formal name for 6 th field hydrologic units.
Surface Fines	Sediments of small size deposited on the surface of a streambed or lake bottom. The upper size threshold for fine sediment for fisheries purposes varies from 0.8 to 605 μ m depending on the observer and methodology used. Results are typically expressed as a percentage of observation points with fine sediment.

Surface Water	All water naturally open to the atmosphere (rivers, lakes, reservoirs, streams, impoundments, seas, estuaries, etc.) and all springs, wells, or other collectors that are directly influenced by surface water.
Suspended Sediments	Fine material (usually sand size or smaller) that remains suspended by turbulence in the water column until deposited in areas of weaker current. These sediments cause turbidity and, when deposited, reduce living space within streambed gravels and can cover fish eggs or alevins.
Taxon	Any formal taxonomic unit or category of organisms (e.g., species, genus, family, order). The plural of taxon is taxa (Armantrout 1998).
Tertiary	An interval of geologic time lasting from 66.4 to 1.6 million years ago. It constitutes the first of two periods of the Cenozoic Era, the second being the Quaternary. The Tertiary has five subdivisions, which from oldest to youngest are the Paleocene, Eocene, Oligocene, Miocene, and Pliocene epochs.
Threatened Species	Species, determined by the U.S. Fish and Wildlife Service, which are likely to become endangered within the foreseeable future throughout all or a significant portion of their range.
Total Maximum Daily Load (TMDL)	A TMDL is a water body's loading capacity after it has been allocated among pollutant sources. It can be expressed on a time basis other than daily if appropriate. Sediment loads, for example, are often calculated on an annual bases. $TMDL = Loading Capacity = Load Allocation + Wasteload Allocation + Margin of Safety$. In common usage, a TMDL also refers to the written document that contains the statement of loads and supporting analyses, often incorporating TMDLs for several water bodies and/or pollutants within a given watershed.
Total Suspended Solids (TSS)	The dry weight of material retained on a filter after filtration. Filter pore size and drying temperature can vary. American Public Health Association Standard Methods (Greenborg, Clescevi, and Eaton 1995) call for using a filter of 2.0 micron or smaller; a 0.45 micron filter is also often used. This method calls for drying at a temperature of 103-105 °C.
Toxic Pollutants	Materials that cause death, disease, or birth defects in organisms that ingest or absorb them. The quantities and exposures necessary to cause these effects can vary widely.
Tributary	A stream feeding into a larger stream or lake.

Trophic State	The level of growth or productivity of a lake as measured by phosphorus content, chlorophyll <i>a</i> concentrations, amount (biomass) of aquatic vegetation, algal abundance, and water clarity.
Turbidity	A measure of the extent to which light passing through water is scattered by fine suspended materials. The effect of turbidity depends on the size of the particles (the finer the particles, the greater the effect per unit weight) and the color of the particles.
Wasteload Allocation (WLA)	The portion of receiving water's loading capacity that is allocated to one of its existing or future point sources of pollution. Wasteload allocations specify how much pollutant each point source may release to a water body.
Water body	A stream, river, lake, estuary, coastline, or other water feature, or portion thereof.
Water Column	Water between the interface with the air at the surface and the interface with the sediment layer at the bottom. The idea derives from a vertical series of measurements (oxygen, temperature, phosphorus) used to characterize water.
Water Pollution	Any alteration of the physical, thermal, chemical, biological, or radioactive properties of any waters of the state, or the discharge of any pollutant into the waters of the state, which will or is likely to create a nuisance or to render such waters harmful, detrimental, or injurious to public health, safety, or welfare; to fish and wildlife; or to domestic, commercial, industrial, recreational, aesthetic, or other beneficial uses.
Water Quality	A term used to describe the biological, chemical, and physical characteristics of water with respect to its suitability for a beneficial use.
Water Quality Criteria	Levels of water quality expected to render a body of water suitable for its designated uses. Criteria are based on specific levels of pollutants that would make the water harmful if used for drinking, swimming, farming, or industrial processes.
Water Quality Limited	A label that describes water bodies for which one or more water quality criterion is not met or beneficial uses are not fully supported. Water quality limited segments may or may not be on a §303(d) list.
Water Quality Limited Segment (WQLS)	Any segment placed on a state's §303(d) list for failure to meet applicable water quality standards, and/or is not expected to meet applicable water quality standards in the period prior to the next list. These segments are also referred to as "§303(d) listed."

Water Quality Modeling	The prediction of the response of some characteristics of lake or stream water based on mathematical relations of input variables such as climate, stream flow, and inflow water quality.
Water Quality Standards	State-adopted and EPA-approved ambient standards for water bodies. The standards prescribe the use of the water body and establish the water quality criteria that must be met to protect designated uses.
Water Table	The upper surface of ground water; below this point, the soil is saturated with water.
Watershed	1) All the land which contributes runoff to a common point in a drainage network, or to a lake outlet. Watersheds are infinitely nested, and any large watershed is composed of smaller “subwatersheds.” 2) The whole geographic region which contributes water to a point of interest in a water body.
Wetland	An area that is at least some of the time saturated by surface or ground water so as to support with vegetation adapted to saturated soil conditions. Examples include swamps, bogs, fens, and marshes.
Young of the Year	Young fish born the year captured, evidence of spawning activity.
